

Effects of the speed of sound at large- N

Ivonne Zavala^{1,*}

¹*Department of Physics, Swansea University, Singleton Park, Swansea, SA2 8PP, UK*

The implications of a smaller than unity speed of sound of the scalar perturbations during cosmic inflation is studied in terms of a model independent large- N approach. We show that the (n_s, r) plane is non-trivially degenerated when the effects of a different than one speed of sound c_s are taken into account. We also discuss the degeneracies due to c_s in the running of the spectral index, α_s . We use the bounds on c_s to constrain such degeneracies.

I. INTRODUCTION

In order to understand the large scale homogeneity and isotropy of the observable universe today, a period of accelerated expansion in the early universe, or cosmic inflation, is needed [1, 2]. Moreover, inflation also provides the seeds of the quantum fluctuations, which gave rise to the large scale structures in the universe that we observe today [3]. Presently, hundreds of different models of inflation driven by a single scalar field exist, with a wide variety of values for the parameters describing the amplitude of the spectra of quantum fluctuations [4].

The stage of inflation and its predictions for the quantum fluctuations, can be described without making precise reference to the microscopic mechanism generating the accelerated expansion [5] (see also [6, 7]). In particular, inflation makes robust predictions for the primordial inhomogeneities. Namely, they are adiabatic, approximately scale invariant and nearly Gaussian. Furthermore, inflation also predicts the existence of primordial gravitational waves.

All these properties are in perfect agreement with current observations. The latest reported values from the Planck collaboration for the inflationary parameters are [8] (at 68% c.l.)

$$n_s - 1 = -0.0397 \pm 0.0073, \quad (1)$$

for the spectral tilt, confirming a percent-level deviation from a scale invariant spectrum, which corresponds to $n_s = 1$. Furthermore, the scale dependence of the spectral index, or the running, defined as $\alpha_s = dn_s/d \ln k$ has been constrained to be

$$\alpha_s = -0.0134 \pm 0.009. \quad (2)$$

In addition, the Planck reported bound on primordial equilateral non-Gaussianity parameter at 68% c.l. is¹ $f_{NL}^{eq} = -42 \pm 75$ [11]. This type of non-Gaussianity results from models where the speed of the inflation perturbations can differ from unity. In this case, the latest

Planck results report a lower bound for this speed of (95% c.l.)

$$c_s \geq 0.02, \quad (3)$$

in an effective field theory parametrisation [12, 13].

Further, the observational upper bound for the ratio of the tensor-to-scalar mode, r , reported by Planck 95% c.l. is

$$r < 0.11, \quad (4)$$

while determination of the primordial origin of the recently reported larger value of $r \sim 0.2$ by the BICEP2 collaboration [14] awaits assessment.

The experimental values for the spectral index and r above, can be neatly described, independently of the underlying model of inflation, when written solely in terms of the number of e-folds N left to the end of inflation after horizon crossing. The number of e-folds is defined as $a = a_f \exp(-N)$, with a the scale factor encoding the accelerated expansion. The relevant interval of N for observations is around $N_* \sim 50 - 60$. For these values, the current data suggest a $1/N_*$ departure from a scale invariant spectrum (1).

We consider the experimental data on n_s and r as an indication to follow a large- N approach to inflation [5, 15–19]. In particular, we take carefully into account the possible degeneracies in the spectral index and tensor-to-scalar ratio due to a smaller than unity speed of sound c_s . A smaller than unity speed of sound for the density perturbations during the inflationary stage, induces primordial non-Gaussianities [20] of equilateral shape and therefore it is constrained by data. Without specifying a concrete model for cosmic inflation, nevertheless, we cannot at present exclude a smaller than light speed for the perturbations.

II. SPEED OF SOUND DEGENERACIES AT LARGE- N

The generic formula for the spectral index n_s in terms of the slow variation parameters, defined as $\epsilon_1 \equiv d \ln H / dN$, $\epsilon_2 \equiv d \ln \epsilon_1 / dN$ and $s \equiv d \ln c_s / dN$, is given by [21]

$$n_s - 1 = -2\epsilon_1 + \epsilon_2 + s, \quad (5)$$

* e.i.zavalacarrasco@swansea.ac.uk

¹ Update: the new constraints in the cosmological parameters above reported by Planck15 [9, 10] are: $n_s = 0.968 \pm 0.006$; $\alpha_s = -0.003 \pm 0.007$; $f_{NL}^{eq} = -16 \pm 70$.

where c_s is the speed of sound, which can be different from unity in general models of cosmic inflation, giving rise to primordial non-Gaussianity of the equilateral shape [20]. The running of the spectral index is further found from its definition, as $\alpha_s = \frac{1}{(\epsilon_1 - 1)} \frac{dn_s}{dN}$. Furthermore, the tensor-to-scalar ratio in the more general case is given by [21]

$$r = 16 c_s \epsilon_1. \quad (6)$$

The current experimental value for the spectral tilt suggests that it should scale as an inverse power of the number of e-folds N at least to leading order in a large- N expansion, independent of the source driving inflation. That is,

$$n_s - 1 = -\frac{A}{N} + \dots \quad (7)$$

with A some constant of order one and the dots referring to subleading terms in a $1/N$ expansion. From (7) and (5), we immediately see that the source of the $1/N$ behaviour of the spectral tilt can originate from any of the slow variation parameters. One can simply assume that only one of them is responsible for such scaling, while the others are subleading. For a unity speed of sound, one can identify two classes of models, according to whether ϵ_1 and ϵ_2 scale both as $1/N$ or only ϵ_2 does [15]. In the present case, one can make a similar classification, including the speed of sound parameter s . Therefore we consider the following expansions²

$$\epsilon_1 = \frac{\alpha}{N^p} \Rightarrow \epsilon_2 = -\frac{p}{N}, \quad s = -\frac{\beta}{N}, \quad (8)$$

where $\alpha > 0$ and β are some constants. This implies that the speed of sound scales with N as

$$c_s = \frac{c_0}{N^\beta}, \quad (9)$$

where c_0 is a positive integration constant, not necessarily of order unity³. Plugging (8) into (5)-(3) we arrive at

$$n_s - 1 = -\frac{2\alpha}{N^p} - \frac{p}{N} - \frac{\beta}{N}, \quad r = \frac{16 c_0 \alpha}{N^{\beta+p}}. \quad (10)$$

Current observations indicate that $A \sim 2$ in (7), which restricts the values of α, β .

We now want to assess the degeneracies in the observational parameters (n_s, r), and the effects on the spectral

tilt running as we allow for a different than unity speed of sound. We do this by changing the values of β , taking into account the experimental bounds on c_s .

Using the current bounds on c_s , we see that for a given value of the constant c_0 , there is a maximum value for the β parameter given by

$$\beta_{max} = \frac{\ln c_0}{\ln N} - \frac{\ln 0.02}{\ln N}, \quad (11)$$

where N is evaluated at horizon crossing, which we take to be $N_* = 60$ for concreteness, but other values can be studied in a similar fashion. Further, the requirement that $c_s \leq 1$, sets a minimum value for β , given c_0 :

$$\beta_{min} = \frac{\ln c_0}{\ln N}. \quad (12)$$

Therefore, for each value of c_0 , there is a range of values for β within the observable window of c_s . We illustrate this in figures 1, where we plot c_s as function of β for different values of c_0 . The allowed range of β set by c_s is evident from this figure.




FIG. 1. Behaviour of the speed of sound as a function of β for different values of c_0 .

The degeneracy sourced by the speed of sound in the spectral tilt and the tensor-to-scalar ratio is illustrated in figure 2 for $\alpha = 0.5$, $p = 1$ and in figure 3 for $\alpha = 0.75$, $p = 2$. In these pictures we see the displacement along c_s , which is reflected in the change of β , for different values of c_0 . The dots correspond to the $(\beta_{min}, \beta_{max})$ range for the given value of c_0 , corresponding to the bounds on $c_s = (1, 0.05)$. As a reference, the chaotic and Starobinsky inflation points are marked with a star and a spade, respectively. These correspond to $\beta_{min} = 0$ ($c_s = 1$), $c_0 = 1$ in the figures. As we can see, the nontrivial effect of the speed of sound, (encoded in β and c_0) is to populate regions in the parameter space (n_s, r) uncovered by the $c_s = 1$ case.

Finally, we show the effect of the speed of sound for the running of the spectral index α_s in figure 4 for $\alpha = 0.5$. As it is evident from (10), variations in c_s move the points

² One could in principle add a third class of models by asking that only $s \sim 1/N$, while ϵ_i are subleading in a large- N expansion. It is not clear if successful models of inflation can have such behaviour, therefore we do not explore this case further.

³ We are considering that scalar and tensor modes cross the horizon at similar instants [21]. One can check that modifications due to this effect [22] are negligible for our discussion. Constraints on the speed of sound taking into account this effect have been studied in [23, 24].




FIG. 2. Speed of sound effects in the $(n_s, \log_{10} r)$ plane for different values of c_0 for $\alpha = 0.5$ and $p = 1$. The dashed curves correspond to the experimental Planck regions (1 and 2σ contours). The dark dots correspond to the allowed range of β set by the bounds on c_s , for the corresponding c_0 . For example, for $c_0 = 1$, $(\beta_{min}, \beta_{max}) = (0, 0.96)$. The red marks correspond to the points of quadratic chaotic (star) and Starobinsky inflation (spades). This plot illustrates the shifts in n_s, r by turning on the speed of sound, populating the intermediate regions between known slow roll models of inflation.

in the (n_s, α_s) plane left upwards. Note however that the shifting is rather mild. From the large- N expression for α_s we see that a similar shifting occurs when varying α and/or p .

As stressed above, we have taken the $1/N$ behaviour of the spectral tilt suggested by observations, without prejudice about the microscopic origin of the stage of inflation and the produced quantum fluctuations. In doing this, one should consider the implications of a different than unity speed of sound for the observable parameters, in particular n_s, r and α_s . Moreover, the latest observational data allow us to use the non-Gaussianity parameters to constrain this variable, as we have discussed.

The important conclusion to draw from this analysis is that, if we do not focus on a particular model for producing the cosmological perturbations, models where some mechanism allows for a non-trivial speed of sound are clearly not excluded by current observations. Moreover, a different than unity speed of sound pushes the tensor-to-scalar ratio and the spectral index to smaller values, covering regions in the (n_s, r) plane, which would seem empty for models with $c_s = 1$. On the other hand, it increases the running of the spectral tilt by only a small amount.




FIG. 3. Similar plot as in figure 2 of the speed of sound effects in the $(n_s, \log_{10} r)$ plane, this time for $\alpha = 0.75$, $p = 2$.

III. DISCUSSION

The current cosmological data indicate a suggestive $1/N$ departure from scale invariance in the spectral tilt when evaluated at the relevant range values of $N = 50 - 60$. Remaining open about the precise origin of the inflationary stage and focusing in a more general description of the parameters in terms of a $1/N$ expansion, one is compelled to include all the small variation parameters contributing to the spectral tilt, (5). In particular, the speed of sound and its variation with N give non-negligible contributions, with nontrivial implications as we have discussed in this note. Variations in the speed of sound source non-Gaussianities, which allow us to constrain the variation on the speed of sound and the degeneracies implied in the (n_s, r) plane and α_s .

We have not been concerned about constraining specific models or classes of models in our analysis. However, there are some single scalar field models of inflation which may be relevant for our discussion. These are k- and DBI-inflation [20, 25, 26], where c_s can become smaller than unity. In consistent single field models of DBI inflation [27], it could be possible to allow for broad ranges of values for β and therefore spectral tilt and tensor-to-scalar ratio, covering a large region in the (n_s, r) plane, unpopulated by $c_s = 1$ models.

In the case of k- and DBI models of inflation, one may also consider the implications for the Lyth bound [28]. In the DBI case, the Lyth bound takes the same form as




FIG. 4. The speed of sound effects in the running of the spectral index for $\alpha = 0.5$. The dashed curves correspond to the allowed Planck regions. The dots correspond to the indicated values of β , with $\beta_{max} = 1.52$ corresponding to $c_0 = 10$. As it is clear from (10), the effect of the speed of sound is to increase by a small amount the value of α_s as we increase β .

in the unity speed of sound case

$$\frac{\Delta\phi}{M_{Pl}} = \int \sqrt{\frac{r}{8}} dN, \quad (13)$$

as show in [29] (here M_{Pl} is the Planck mass). However,

since the speed of sound now enters into the tensor-to-scalar ratio (see (6)), the associated field range will be smaller as compared to the unity speed of sound case. One could nevertheless study the implications of c_s for the field range along the lines of [17, 18] in a large- N approach.

In summary, we have analysed the implications of including the speed of sound in the general formulae for the observable inflationary parameters, namely the spectral tilt, and its running, as well as the tensor-to-scalar ratio, independent of the mechanism producing the inflationary stage in the early universe. We have done so by writing all the slow variation parameters in a $1/N$ expansion in the number of e-folds, as suggested by current observations. We have used the current constraints on the relevant non-Gaussianity parameter sourced by c_s to explore and determine the implied degeneracies in the observable parameters. As we have seen, non-trivial motion in the (n_s, r) plane can occur towards smaller values of n_s and r , covering parts of the parameter space, unpopulated by the $c_s = 1$ case.

A possible way to break the degeneracies implied by the speed of sound, could come in the future from a scale dependence of the non-Gaussianity parameter, if non-Gaussianity is detected [30, 31].

It would also be interesting to perform a model independent analysis as has been done in [6, 7] taking into account the small variation parameter s .

ACKNOWLEDGMENTS

I would like to thank Diederik Roest, Marco Scalisi and Gianmassimo Tasinato for useful comments on the manuscript.

-
- [1] A. H. Guth, “The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems”, *Phys.Rev.* **D23** (1981) 347–356.
 - [2] A. D. Linde, “A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems”, *Phys.Lett.* **B108** (1982) 389–393.
 - [3] V. F. Mukhanov and G. V. Chibisov, “Quantum Fluctuation and Nonsingular Universe. (In Russian)”, *JETP Lett.* **33** (1981) 532–535.
 - [4] J. Martin, C. Ringeval, and V. Vennin, “Encyclopedia Inflationaris”, *Phys.Dark Univ.* (2014), [arXiv:1303.3787 \[astro-ph.CO\]](#).
 - [5] V. Mukhanov, “Quantum Cosmological Perturbations: Predictions and Observations”, *Eur.Phys.J.* **C73** (2013) 2486, [arXiv:1303.3925 \[astro-ph.CO\]](#).
 - [6] L. Barranco, L. Boubekeur, and O. Mena, “A model-independent fit to Planck and BICEP2 data”, *Phys.Rev.* **D90** (2014) 063007, [arXiv:1405.7188 \[astro-ph.CO\]](#).
 - [7] L. Boubekeur, E. Giusarma, O. Mena, and H. Ramírez, “Phenomenological approaches of inflation and their equivalence”, [arXiv:1411.7237 \[astro-ph.CO\]](#).
 - [8] **Planck Collaboration** Collaboration, P. Ade *et al.*, “Planck 2013 results. XXII. Constraints on inflation”, *Astron.Astrophys.* **571** (2014) A22, [arXiv:1303.5082 \[astro-ph.CO\]](#).
 - [9] **Planck Collaboration** Collaboration, P. Ade *et al.*, “Planck 2015. XX. Constraints on inflation”, [arXiv:1502.02114 \[astro-ph.CO\]](#).
 - [10] **Planck Collaboration** Collaboration, P. Ade *et al.*, “Planck 2015 results. XVII. Constraints on primordial non-Gaussianity”, [arXiv:1502.01592 \[astro-ph.CO\]](#).
 - [11] **Planck Collaboration** Collaboration, P. Ade *et al.*, “Planck 2013 Results. XXIV. Constraints on primordial non-Gaussianity”, *Astron.Astrophys.* **571** (2014) A24, [arXiv:1303.5084 \[astro-ph.CO\]](#).
 - [12] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan, and L. Senatore, “The Effective Field Theory of Inflation”, *JHEP* **0803** (2008) 014, [arXiv:0709.0293](#)

- [hep-th].
- [13] S. Weinberg, “Effective Field Theory for Inflation”, *Phys.Rev.* **D77** (2008) 123541, [arXiv:0804.4291 \[hep-th\]](#).
 - [14] **BICEP2 Collaboration** Collaboration, P. Ade *et al.*, “Detection of B-Mode Polarization at Degree Angular Scales by BICEP2”, *Phys.Rev.Lett.* **112** (2014) 241101, [arXiv:1403.3985 \[astro-ph.CO\]](#).
 - [15] D. Roest, “Universality classes of inflation”, *JCAP* **1401** (2014) no. 01, 007, [arXiv:1309.1285 \[hep-th\]](#).
 - [16] J. Garcia-Bellido and D. Roest, “Large- N running of the spectral index of inflation”, *Phys.Rev.* **D89** (2014) no. 10, 103527, [arXiv:1402.2059 \[astro-ph.CO\]](#).
 - [17] J. Garcia-Bellido, D. Roest, M. Scalisi, and I. Zavala, “Can CMB data constrain the inflationary field range?”, *JCAP* **1409** (2014) 006, [arXiv:1405.7399 \[hep-th\]](#).
 - [18] J. Garcia-Bellido, D. Roest, M. Scalisi, and I. Zavala, “Lyth bound of inflation with a tilt”, *Phys.Rev.* **D90** (2014) no. 12, 123539, [arXiv:1408.6839 \[hep-th\]](#).
 - [19] P. Creminelli, S. Dubovsky, D. L. Nacir, M. Simonovi?, G. Trevisan, *et al.*, “Implications of the scalar tilt for the tensor-to-scalar ratio”, [arXiv:1412.0678 \[astro-ph.CO\]](#).
 - [20] M. Alishahiha, E. Silverstein, and D. Tong, “DBI in the sky”, *Phys.Rev.* **D70** (2004) 123505, [arXiv:hep-th/0404084 \[hep-th\]](#).
 - [21] J. Garriga and V. F. Mukhanov, “Perturbations in k-inflation”, *Phys.Lett.* **B458** (1999) 219–225, [arXiv:hep-th/9904176 \[hep-th\]](#).
 - [22] N. Agarwal and R. Bean, “Cosmological constraints on general, single field inflation”, *Phys.Rev.* **D79** (2009) 023503, [arXiv:0809.2798 \[astro-ph\]](#).
 - [23] D. Baumann, D. Green, and R. A. Porto, “B-modes and the Nature of Inflation”, *JCAP* **1501** (2015) no. 01, 016, [arXiv:1407.2621 \[hep-th\]](#).
 - [24] G. A. Palma and A. Soto, “B-modes and the sound speed of primordial fluctuations”, [arXiv:1412.0033 \[hep-th\]](#).
 - [25] C. Armendariz-Picon, T. Damour, and V. F. Mukhanov, “k - inflation”, *Phys.Lett.* **B458** (1999) 209–218, [arXiv:hep-th/9904075 \[hep-th\]](#).
 - [26] E. Silverstein and D. Tong, “Scalar speed limits and cosmology: Acceleration from D-celeration”, *Phys.Rev.* **D70** (2004) 103505, [arXiv:hep-th/0310221 \[hep-th\]](#).
 - [27] A. Avgoustidis and I. Zavala, “Warped Wilson Line DBI Inflation”, *JCAP* **0901** (2009) 045, [arXiv:0810.5001 \[hep-th\]](#).
 - [28] D. H. Lyth, “What would we learn by detecting a gravitational wave signal in the cosmic microwave background anisotropy?”, *Phys.Rev.Lett.* **78** (1997) 1861–1863, [arXiv:hep-ph/9606387 \[hep-ph\]](#).
 - [29] D. Baumann and L. McAllister, “A Microscopic Limit on Gravitational Waves from D-brane Inflation”, *Phys.Rev.* **D75** (2007) 123508, [arXiv:hep-th/0610285 \[hep-th\]](#).
 - [30] X. Chen, “Running non-Gaussianities in DBI inflation”, *Phys.Rev.* **D72** (2005) 123518, [arXiv:astro-ph/0507053 \[astro-ph\]](#).
 - [31] C. T. Byrnes, S. Nurmi, G. Tasinato, and D. Wands, “Scale dependence of local fNL”, *JCAP* **1002** (2010) 034, [arXiv:0911.2780 \[astro-ph.CO\]](#).